

# Injectables

## Product Catalogue


Prevention & Treatment of Diseases  
and Infections in Livestock



enriches the agriculture industry by providing adequate and quality nutritional solutions.

**BETTER INJECTABLES,  
BETTER ANIMAL HEALTH.**

[HIPRO-FEED.COM](http://HIPRO-FEED.COM)





additives that guarantee the best protection of your investment, leading to a maximum profitability.

#### **Vision**

To help raise healthy animals by supplying premium quality animal nutrition products.

#### **Mission**

Prioritize the growth, performance and health needs of our clients animals.

#### **Objectives**

- Continually analyze the best solutions to improve our range of products.
- Establish and build strong relation with our existing and new clients.



03 – Protyl-20



04 – Moxpro15 LA



05 – Gentapro



06 – Prophos12



15 – Prociprox10



16 – Provaquone



17 – Pro-E+S



18 – Protil30



07 – Proflor30



08 – Proclosan



09 – Oxypro20 LA



10 – Oxypro30 LA



19 – Pronalgin



20 – Prodexam



21 – Proenrox



22 – Prosultrim



Usage: Subcutaneous administration.

Livestock (sheep, goat, cattle): 1 mL per 50 kg of body weight.

#### Packing

Promectin 1% is packed in 100mL bottles.

#### Safety Precautions!

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

SUBCUTANEOUS ADMINISTRATION METHOD  
FOR LIVESTOCK.



Usage: Subcutaneous administration.

Livestock (sheep, goat, cattle): 1 mL per 50 kg of body weight.

#### **Packing**

Promectin 2% is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

SUBCUTANEOUS ADMINISTRATION METHOD  
FOR LIVESTOCK.



#### Dosage

Usage: Intramuscular administration.

Livestock (sheep, goat, cattle): 1 mL per 20 Kg body weight for 3-5 days.

#### Packing

Protyl-20 is packed in 100mL bottles.

#### Safety Precautions!

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

#### INTRAMUSCULAR ADMINISTRATION METHOD FOR LIVESTOCK.



#### Dosage

Usage: Intramuscular administration.

Livestock (sheep, goat, cattle): 1 mL per 10 kg body weight, repeatable if necessary after 48 hours.

#### Packing

Moxpro-15 LA is packed in 100mL bottles.

#### Safety Precautions!

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

#### INTRAMUSCULAR ADMINISTRATION METHOD FOR LIVESTOCK.



#### **Dosage**

Usage: Intramuscular administration.

Livestock (sheep, goat, cattle): Twice daily 1 mL per 20 - 40 Kg body weight for 3 days.

#### **Packing**

Gentapro is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR ADMINISTRATION METHOD  
FOR LIVESTOCK.



a support of muscular physiology as an effective tonic in cases of stress, over-exertion, exhaustion and reduced resistance.

#### **Dosage**

Usage: Intravenous, Intramuscular or Subcutaneous administration.  
Cattle and horses: 10 mL per animal (intravenous). Sheep and goats: 5 mL per animal.

#### **Packing**

Prophos12 is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAVENOUS, INTRAMUSCULAR OR SUBCUTANEOUS  
ADMINISTRATION METHOD FOR LIVESTOCK.



Usage: Subcutaneous or intramuscular administration.  
Livestock (sheep, goat, cattle): 1 mL per 20 Kg body weight for 3-5 days.

#### **Packing**

Proflo30 is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR OR SUBCUTANEOUS ADMINISTRATION  
METHOD FOR LIVESTOCK.



Usage: Subcutaneous administration.

Livestock (sheep, goat, cattle): 1 mL per 20-40 Kg body weight for 3-5 days.

#### **Packing**

Proclosan is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

SUBCUTANEOUS ADMINISTRATION METHOD  
FOR LIVESTOCK.



Livestock (sheep, goat, cattle): General: 1 mL per 10 Kg body weight. This dosage can be repeated after 48 hours when necessary.  
*Do not administer more than 20 mL in cattle and more than 5 mL in calves, goats and sheep per injection site.*

#### **Packing**

Oxypro20 LA is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR OR SUBCUTANEOUS  
ADMINISTRATION METHOD FOR LIVESTOCK.



Livestock (sheep, goat, cattle): General: 1 mL per 20-30 Kg body weight. This dosage can be repeated after 48 hours when necessary.  
*Do not administer more than 20 mL in cattle and more than 5 mL in calves, goats and sheep per injection site.*

#### **Packing**

Oxypro30 LA is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR OR SUBCUTANEOUS  
ADMINISTRATION METHOD FOR LIVESTOCK.



#### Dosage

Usage: Subcutaneous or intramuscular administration.

Livestock (sheep, goat, cattle): 1 mL per 10 Kg body weight for 3-5 days.

#### Packing

Proiro40 is packed in 100mL bottles.

#### Safety Precautions!

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR OR SUBCUTANEOUS  
ADMINISTRATION METHOD FOR LIVESTOCK.



#### **Dosage**

Usage: Intramuscular administration.

Livestock (sheep, goat, cattle): 1 mL per 20 Kg body weight for 3-5 days.

#### **Packing**

Prospeclin is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR ADMINISTRATION METHOD  
FOR LIVESTOCK.



#### Dosage

Usage: Intramuscular administration.

Livestock (sheep, goat, cattle): 1 mL per 10-20 Kg body weight for 3 days.

*Shake well before use and do not administer more than 20 mL in cattle and more than 5 mL in calves, sheep and goats per injection site.*

#### Packing

Prostrept400 is packed in 100mL bottles.

#### Safety Precautions!

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

#### INTRAMUSCULAR ADMINISTRATION METHOD FOR LIVESTOCK.



Usage: Subcutaneous or intramuscular administration.  
Livestock (sheep, goat, cattle): 1 mL per 5 Kg body weight for 3-5 days.

#### Packing

Prosulfa-333 is packed in 100mL bottles.

#### Safety Precautions!

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

SUBCUTANEOUS OR INTRAMUSCULAR  
ADMINISTRATION METHOD FOR LIVESTOCK.

**Dosage**

Usage: Intravenous or intramuscular administration.

Livestock (sheep, goat, cattle): Twice daily 5 mL per 75-150 Kg body weight for 3-5 days.

**Packing**

Prociprox10 is packed in 100mL bottles.

**Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAVENOUS OR INTRAMUSCULAR  
ADMINISTRATION METHOD FOR LIVESTOCK.



#### **Dosage**

Usage: Intramuscular administration.

Livestock (sheep, goat, cattle): 1 mL per 20 Kg body weight. In severe cases the treatment may be repeated within 48-72 hours.

#### **Packing**

Provaquone is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

#### **INTRAMUSCULAR ADMINISTRATION METHOD FOR LIVESTOCK.**



destroying oxidising agents.

#### **Dosage**

Usage: Intramuscular or subcutaneous administration.  
Livestock (sheep, goat, cattle): 2 mL per 10 Kg body weight.

#### **Packing**

Pro-E+S is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR OR SUBCUTANEOUS  
ADMINISTRATION METHOD FOR LIVESTOCK.

**Dosage**

Usage: Subcutaneous administration.  
Livestock (sheep, goat, cattle): 1 mL per 30 Kg body weight.

**Packing**

Protil30 is packed in 100mL bottles.

**Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

SUBCUTANEOUS ADMINISTRATION METHOD  
FOR LIVESTOCK.



#### **Dosage**

Usage: Intramuscular or intravenous administration.  
Livestock (sheep, goat, cattle): 2 mL per 50 Kg body weight.

#### **Packing**

Pronalgin is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR OR INTRAVENOUS  
ADMINISTRATION METHOD FOR LIVESTOCK.

**Dosage**

Usage: Intramuscular or intravenous administration.  
Livestock (sheep, goat, cattle): 2 mL per 50 Kg body weight.

**Packing**

Prodexam is packed in 100mL bottles.

**Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR OR INTRAVENOUS  
ADMINISTRATION METHOD FOR LIVESTOCK.

**Dosage**

Usage: Intramuscular or subcutaneous administration.

Livestock (sheep, goat, cattle): 1 mL per 20-40 Kg body weight for 3-5 days.

**Packing**

Proenrox is packed in 100mL bottles.

**Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

INTRAMUSCULAR OR SUBCUTANEOUS  
ADMINISTRATION METHOD FOR LIVESTOCK.



calves, cattle, goats and sheep.

#### **Dosage**

Usage: Intramuscular administration.

Livestock (sheep, goat, cattle): 1 mL per 10 Kg body weight for 3-5 days.

#### **Packing**

Prosultrim is packed in 100mL bottles.

#### **Safety Precautions!**

1. Store in a dry place below 30°C.
2. Keep out of reach of children.
3. Strictly for animal use only.
4. Close tightly after every usage.

#### INTRAMUSCULAR ADMINISTRATION METHOD FOR LIVESTOCK.



DYNAMIC.  
EXCELLENCE.  
CUTTING-EDGE.  
HIPRO.



**HIPRO İÇ VE DIŐ TİCARET SAN.VE TİC. LTD. ŐTİ**

Zafer Sb Mah. Ayhan Sok. S-4 Binasi No:10/2  
Ege Serbest Bolgesi, Gaziemir - Izmir, Turkey

**[HIPRO-FEED.COM](http://HIPRO-FEED.COM)**

